

INTRAVENOUS (IV) OXYTOCINE REGIMEN FILE

Patient name	Patient #	Bed #
Indication	Date	
Bishop score		

PROTOCOL

Dilute 5 IU in 500 ml or 10 IU in 1 litre of Ringer lactate or 0.9% sodium chloride solution.

Start at 5 to 8 drops / minute, then increase by 5 to 8 drops / minute every 30 minutes, until contractions are effective (3 contractions of more than 40 seconds in 10 minutes). On average, 20 drops / minute result in satisfactory uterine contractions. Do not exceed 60 drops / minute.

In case of previous C-section (low transverse incision), grand multiparity and/or overdistended uterus the maximum infusion rate of 30 drops / minute.

In case of fœtal distress, uterine hyperkinesia (more than 5 contractions in 10 minutes), or uterine hypertony (absence of uterine relaxation): STOP the oxytocin.

Every 30 minutes monitor: Close maternal monitoring (check for hyperstimulation, imminent rupture and dystocia)

Close fœtal monitoring (chek for decelerations in heart rate after the contractions)

IV dri	p rate		Contractions		Foetal heart rate		
hour	drops/ minute	yes/no	/ 10 minutes	seconds	/ minute	Remarks	Signature

Other medication given:		